

## Recombinant Human FLT3LG (C-6His)

Catalog#:AC13161 Derived from Human Cells

<b>DESCRIPTION</b>	<p>Recombinant Human Fms-like Tyrosine Kinase 3 Ligand is produced by our Mammalian expression system and the target gene encoding Thr27-Pro184 is expressed with a 6His tag at the C-terminus.</p> <p>Accession#: P49771</p> <p>Known as: Fms-Related Tyrosine Kinase 3 Ligand; Flt3 Ligand; Flt3L; SL Cytokine; FLT3LG</p>
<b>FORMULATION</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
<b>SHIPPING</b>	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
<b>STORAGE</b>	<p>Lyophilized protein should be stored at &lt; -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at &lt; -20°C for 3 months.</p>
<b>RECONSTITUTION</b>	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100µg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
<b>QUALITY CONTROL</b>	<p>Mol Mass: 19kDa AP Mol Mass: 24-32kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test.</p>
<b>BACKGROUND</b>	<p>Fms-Related Tyrosine Kinase 3 Ligand (FLT3LG) is a hematopoietic four helical bundle cytokine. Mature human Flt-3 Ligand consists of an extracellular domain (ECD) with a cytokine-like domain and a juxtamembrane tether region, a transmembrane segment, and a cytoplasmic tail. Human and mouse Flt-3 Ligand show cross-species activity. Flt-3 Ligand is expressed as a noncovalently-linked dimer by T cells and bone marrow and thymic fibroblasts. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors by activation of Flt 3 receptor.</p>
<b>SDS-PAGE</b>	