

Recombinant Mouse OX40L (N-8His) Catalog#:AC13252 Derived from Human Cells

DESCRIPTION	Recombinant Mouse OX40 Ligand is produced by our Mammalian expression system and the target gene encoding Ser51-Leu198 is expressed with a 8His tag at the N-terminus. Accession#:P43488 Known as:Tumor necrosis factor ligand superfamily member 4; OX40 ligand; OX40L; CD252; Tnfsf4
FORMULATION	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
SHIPPING	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY CONTROL	Mol Mass: 17.7kDa AP Mol Mass: 20-23 kDa, reducing conditions. Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test.
BACKGROUND	OX40 ligand (OX40L), also called CD252, is a single-pass type II membrane protein of the TNF/TNF receptor superfamily. OX40L is expressed by DCs, macrophages and B cells and signals via its cognate receptor OX40 which is mainly expressed on APCs. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells. OX40L–OX40 co-stimulation leads to activation of TNF receptor associated factor (TRAF) 2, 3 and 5. This pathway has been shown to prolong the survival of effector CD4+Th cells as well as contributes to generation of memory T cells.
	kDa MK R 120 90 60 90 60 40 40 40 40 20 14 14