

Anti-MBP Polyclonal Antibody

Cat: AC51675


Summary:

- | | |
|---|---|
| 【Product name】 : Anti-MBP antibody | 【Source】 : Rabbit |
| 【Isotype】 : IgG | 【Species reactivity】 : Human Mouse Rat |
| 【Swiss Prot】 : P02686 | 【Gene ID】 : 4155 |
| 【Calculated】 : MW:33kDa | 【Observed】 : MW:33kDa |
| 【Purification】 : Affinity purification | |
| 【Tested applications】 : WB IHC IF | |
| 【Recommended dilution】 : WB 1:500-2000. IHC 1:50-200. IF 1:50-200. | |
| 【WB Positive sample】 : HL-60 | |
| 【IHC Positive sample】 : Rat kidney tissue | |
| 【Subcellular location】 : Cell membrane Membrane Nucleus | |
| 【Immunogen】 : Recombinant protein of human MBP | |
| 【Storage】 : Shipped at 4°C. Upon delivery aliquot and store at -20°C | |


Background:

The protein encoded by the classic MBP gene is a major constituent of the myelin sheath of oligodendrocytes and Schwann cells in the nervous system. However, MBP-related transcripts are also present in the bone marrow and the immune system. These mRNAs arise from the long MBP gene (otherwise called "Golli-MBP") that contains 3 additional exons located upstream of the classic MBP exons. Alternative splicing from the Golli and the MBP transcription start sites gives rise to 2 sets of MBP-related transcripts and gene products. The Golli mRNAs contain 3 exons unique to Golli-MBP, spliced in-frame to 1 or more MBP exons. They encode hybrid proteins that have N-terminal Golli aa sequence linked to MBP aa sequence. The second family of transcripts contain only MBP exons and produce the well characterized myelin basic proteins. This complex gene structure is conserved among species suggesting that the MBP transcription unit is an integral part of the Golli transcription unit and that this arrangement is important for the function and/or regulation of these genes.

Verified picture



Western blot analysis with MBP antibody diluted at 1:1000



Immunohistochemistry of paraffin-embedded Rat kidney tissue with MBP antibody diluted at 1:200