

Recombinant Human TREM-1 (C-6His)

Catalog#:AC13132 Derived from Human Cells

DESCRIPTION	Recombinant Human Triggering Receptor Expressed on Myeloid Cells 1 is produced by our Mammalian expression system and the target gene encoding Ala21-Arg200 is expressed with a 6His tag at the Cterminus. Accession#: Q9NP99 Known as: Triggering Receptor Expressed on Myeloid Cells 1; TREM-1; Triggering Receptor Expressed on Monocytes 1; CD354; TREM1
FORMULATION	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
SHIPPING	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY CONTROL	Mol Mass: 21.3kDa AP Mol Mass: 32-40kDa, reducing conditions. Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) is a transmembrane protein with a single Ig-like domain. TREM-1 associates with the adapter protein, DAP12, to deliver an activating signal. TREM-1 is expressed on blood neutrophils and monocytes, and the expression is up-regulated by bacterial LPS. TREM1 is expressed at high levels on neutrophils of patients with microbial sepsis and in mice with a TREM-1/Fc fusion protein protected mice against LPS-induced shock. Human TREM-1 shares 42% sequence homology with mouse TREM-1.
SDS-PAGE 20	