


Recombinant Mouse SEMA4A (C-6His)

Catalog#:AC13213 Derived from Human Cells

DESCRIPTION	<p>Recombinant Mouse T-cell-specific surface glycoprotein CD28 is produced by our Mammalian expression system and the target gene encoding Asn20-Lys149 is expressed with a Fc, 6His tag at the C-terminus.</p> <p>Accession#: P31041</p> <p>Known as: T-cell-specific surface glycoprotein CD28;CD28</p>
FORMULATION	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100µg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass: 43kDa AP Mol Mass: 60kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test.</p>
BACKGROUND	<p>T-cell-specific surface glycoprotein CD28, contains an Ig-like V-type domain. CD28 is one of the proteins expressed on T cells that provide co-stimulatory signals, which are required for their activation. It is the receptor for CD80 and CD86. When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells. It involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival.</p>
 <p>SDS-PAGE</p> <p>kDa MK R</p> <p>120</p> <p>90</p> <p>60</p> <p>40</p> <p>30</p> <p>20</p> <p>14</p>	