


Recombinant Mouse IL-23R (C-Fc)

Catalog#:AC13246 Derived from Human Cells

DESCRIPTION	<p>Recombinant Mouse Interleukin-23 Receptor is produced by our Mammalian expression system and the target gene encoding Gly24-Asp372 is expressed with a Fc tag at the C-terminus.</p> <p>Accession#:Q8K4B4</p> <p>Known as: Interleukin-23 receptor; IL23r; IL-23 receptor; IL-23R; interleukin 23 receptor</p>
FORMULATION	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100μg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass: 67.4kDa AP Mol Mass: 90-110kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>Interleukin 23 receptor (IL23R), a heterodimer of the IL12 receptor β1 (IL12Rβ1) and IL12Rβ2, is a type I cytokine receptor for IL23. IL23R is comprised of two subunits, the IL12Rβ1 subunit, which is shared with several cytokines, and a subunit that is unique to IL-23. IL23, after binding to IL23R, activates memory T cells and mediates pro-inflammatory activities in part by the production of IL17 through activation of TH17 lymphocytes. IL23R is expressed on T cells, NK cells, dendritic cells, and macrophages. In fact, polymorphisms of the IL23R gene were reported to be associated with susceptibility to inflammatory diseases and autoimmune diseases such as psoriasis, multiple sclerosis, Graves's ophthalmopathy and inflammatory bowel diseases. The IL23R is known to be critically involved in the carcinogenesis of different malignant tumor.</p>
 <p>SDS-PAGE</p> <p>kDa MK R</p> <p>120</p> <p>90</p> <p>60</p> <p>40</p> <p>30</p>	