

## Recombinant Human IL-35 (C-Fc)

Catalog#:AC13257 Derived from Human Cells

DESCRIPTION	Recombinant Human Interleukin-35 is produced by our Mammalian expression system and the target gene encoding Arg23-Ser219&Arg21-Lys229 is expressed with a Fc tag at the C-terminus.  Accession#:P29459&Q14213  Known as:Interleukin 35; IL35; hIL35
FORMULATION	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
SHIPPING	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.  Reconstituted protein solution can be stored at 4-7°C for 2-7 days.  Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY CONTROL	Mol Mass: 73.7kDa AP Mol Mass: 80-120kDa, reducing conditions. Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	Interleukin-35 is a newest and potent suppressive cytokine of the IL-12 family. Unlike other members of the IL-12 family which are produced mainly by antigen-presenting cells (APCs), IL-35 has been identified as a potentially immunosuppressive cytokine primarily produced by regulatory T (Treg) cells, and to a lesser extent by activated dendritic cells and macrophages. IL-35 is composed of the IL-12 p35 subunit and the IL27 Epstein – Barr virus-induced protein 3 (EBI3) subunit. IL-35 plays an important role in many disease models and has been recently shown to have additional functions aside from inhibition of proliferation, including inducing its own expression in non-Treg cells.
	SDS-PAGE    KDa   MK   R     120