

Glutathione Reductase from baker's yeast

Product Number: G79690

Enzymatic Activity: 100-300 units/mg protein

Storage Temperature: 2-8 ° C

Product Description

Enzyme Commission (EC) Number: 1.6.4.2

CAS Number: 9001-48-3

Molecular Weight: 124 kDa

Glutathione reductase from Baker's yeast is a flavoprotein homodimer consisting of two equal subunits. Each subunit has one mole of FAD which is noncovalently bound. Glutathione reductase is also a sulfhydryl protein containing a total of six sulfhydryl groups. The enzyme catalyzes the following reaction:

Glutathione(Oxidized) + β -NADPH $\rightarrow \beta$ -NADP + 2 Glutathione(Reduced)

The Km values for the enzyme are: oxidized glutathione (61 μ M) and β -NADPH (7.6 μ m).

Gltathione reductase is inhibited by the following inhibitors:

N-alkylmaleimides; benzylselenosulphate; 2-chloroethylisocyanate; Cu²⁺; 2,4-dihydroxybenzylamine;

1-fluoro-2,4-dinitrobenzene; p-nitrobenzylselenosulphate; 2-triazine-5-nitrofuran

Unit Definition

One unit will reduce 1.0 µmole of oxidized glutathione per min at pH 7.6 at 25 °C. Physical form

Suspension in 3.6 M (NH4)2SO4, pH 7.0, containing 0.1 mM dithiothreitol

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

产品仅供科研!



Storage/Stability

This product is offered as an ammonium sulfate suspension. Diluted stock solutions should not be prepared.

